

Cystitis (HPC)

1. Yun² Bai², Li³ Bai², Tong¹ Shen⁴, and Shen⁴ Guan¹ – treat every other day and symptoms should be completely resolved within 10 days (HPC):

- **Yun² Bai²:** This point is found 2 *cun* obliquely anterior and distal to Bei¹ Mian⁴. Needle 0.3 to 0.5 *cun*. {Six Bowels and Lung Auxiliary Branch Reaction Areas} (Bei¹ Mian⁴ is found in the large depression immediately lateral (distal) to the acromion process of the scapula when the arm is abducted to 90°.)
- **Li³ Bai²:** This point is located 2 *cun* distal and slightly dorsal to Yun² Bai². Needle 0.3 to 0.5 *cun*. {Kidney Auxiliary and Lung Branch Reaction Areas}
- **Tong¹ Shen⁴:** This point is located in the depression found at the superomedial angle of the patella. Needle 0.3 to 0.5 *cun*. {Kidney Reaction Area}
- **Shen⁴ Guan¹:** This point lies 1.5 *cun* immediately distal to Tian¹ Huang². Needle 0.5 to 1.0 *cun*. (Tian¹ Huang² is located in the depression found immediately distal to the medial tibial condyle, and lies 2.5 *cun* distal to the articular surface of the medial tibial condyle.) {Six Bowels Reaction Area}

