

Erectile Dysfunction (HPC)

1. Shen⁴ Guan¹, Di⁴ Huang², Ren² Huang², Tong¹ Shen⁴, and Tong¹ Wei⁴ (HPC):

- **Shen⁴ Guan¹**: This point lies 1.5 *cun* immediately distal to Tian¹ Huang². Needle 0.5 to 1.0 *cun*. (Tian¹ Huang² is located in the depression found immediately distal to the medial tibial condyle, and lies 2.5 *cun* distal to the articular surface of the medial tibial condyle.) {Six Bowels Reaction Area}
- **Di⁴ Huang²**: This point is located 7 *cun* proximal to the medial malleolus along the medial border of the tibia. Needle inferior to superior at a 45⁰ angle 1.0 to 1.8 *cun*. {Kidney Branch Reaction Area}
- **Ren² Huang²**: This point is located 3 *cun* proximal to the medial malleolus along the medial border of the tibia. Needle 0.6 to 1.2 *cun*. {Kidney Branch Reaction Area}
- **Tong¹ Shen⁴**: This point is located in the depression found at the superomedial angle of the patella. Needle 0.3 to 0.5 *cun*. {Kidney Reaction Area}
- **Tong¹ Wei⁴**: This point is located 2.0 *cun* immediately proximal to Tong¹ Shen⁴. Needle 0.5 to 1.0 *cun*. {Kidney Reaction Area} (Tong¹ Shen⁴ is located in the depression found at the superomedial angle of the patella)

