

Glomerulonephritis (LCH-1)

1. Wan⁴ Shun⁴ Yi¹, Wan⁴ Shun⁴ Er⁴, Shui³ Xiang⁴, and Shui³ Xian¹ (LCH):

- **Wan⁴ Shun⁴ Yi¹**: This point is located 2.5 *cun* distal to the anterior distal transverse wrist crease, anterior to the 5th metacarpal bone. Needle 0.5 to 1.5 *cun*. {Kidney Branch Reaction Area}
- **Wan⁴ Shun⁴ Er⁴**: This point is located 1.5 *cun* distal to the anterior distal transverse wrist crease, anterior to the 5th metacarpal bone. Needle 0.2 to 0.4 *cun*. {Kidney Branch Reaction Area}
- **Shui³ Xiang⁴**: This point is located in the depression immediately anterior to the Achilles tendon, 2.0 *cun* posterior to the apex (as viewed from posterior to anterior) of the medial malleolus, on the medial side of the ankle. Needle 0.3 to 0.5 *cun* – one may even needle to penetrate through to the lateral side along the anterior border of the Achilles tendon. {Kidney Branch and Brain Reaction Areas}
- **Shui³ Xian¹**: This point is located 2.0 *cun* distal to Shui³ Xiang⁴. It may be found by coursing 2.0 *cun* posterior to the apex (as viewed from posterior to anterior) of the medial malleolus, and then 2.0 *cun* immediately distal. Needle 0.5 *cun*. {Kidney Branch and Brain Reaction Areas}

