

Orchitis (HPC-1)

1. Da⁴ Jian⁴, Xiao³ Jian⁴, Wai⁴ Jian⁴, Tian¹ Huang², and Ren² Huang² – treat daily until symptoms resolve (HPC):

- **Da⁴ Jian⁴:** This point is located on the anterior surface of the proximal phalanx of the index finger, 0.3 *cun* radial (lateral) to the exact center of said phalanx. Needle 0.2 to 0.3 *cun*. *BILATERAL NEEDLING CONTRAINDICATED*. {Heart Organ and Six Bowels Branch Reaction Areas} (This point is located at the proximal-to-distal center of digit line B; to locate the center of digit line B, simply bisect the distance from the anterior metacarpophalangeal skin crease to the anterior proximal interphalangeal skin crease along the course of digit line B.)
- **Xiao³ Jian⁴:** This point is located 0.2 *cun* distal to Da⁴ Jian⁴. Needle 0.2 to 0.3 *cun*. *BILATERAL NEEDLING CONTRAINDICATED*. {Lung Branch, Heart Organ, and Six Bowels Branch Reaction Areas}
- **Wai⁴ Jian⁴:** This point is located on the anterior surface of the middle phalanx of the index finger, 0.2 *cun* radial (lateral) to the median plane of said phalanx, and one third the distance from the proximal interphalangeal skin crease to the distal interphalangeal skin crease. Needle 0.2 *cun*. *BILATERAL NEEDLING CONTRAINDICATED*. {Heart Organ and Six Bowels Branch Reaction Areas} (This point is located one third the distance from the proximal interphalangeal skin crease to the distal interphalangeal skin crease, along the course of digit line B.)
- **Tian¹ Huang²:** This point is found in the depression immediately distal to the medial tibial condyle, and lies 2.5 *cun* distal to the articular surface of the medial tibial condyle. Needle 0.5 to 1.0 *cun*. *MOXABUSTION CONTRAINDICATED*. {Kidney, Six Bowels, and Heart Branch Reaction Areas}
- **Ren² Huang²:** This point is located 3 *cun* proximal to the medial malleolus along the medial border of the tibia. Needle 0.6 to 1.2 *cun*. {Kidney Branch Reaction Area}

