



Paresthesiae: Pedal (LKC-2)

1. Needle Li³ Bai², Yun² Bai², ‘let’ Wu³ Ling³, San¹ Jiang¹, and the affected foot (LKC):




- **Li³ Bai²:** This point is located 2.0 *cun* anterior to Jian⁴ Zhong¹, on the same transverse plane; Needle perpendicular insertion 0.5 to 1.5 *cun* or oblique insertion from distal to proximal 1.5 *cun* to 2.0 *cun*. {Lung Branch and Kidney Auxiliary Reaction Areas} (Jian⁴ Zhong¹ is located on the lateral aspect of the proximal brachium 4.5 *cun* distal to LI-15^{WHO} on the Hand Yang² Ming² Large Intestine Channel.)
- **Yun² Bai²:** This point is located on the anterolateral aspect of the proximal brachium, 2.0 *cun* immediately anterior to, and on the same transverse plane as, Jian¹ Zhong¹; Needle perpendicularly 1.0 to 2.0 *cun* or oblique insertion, from proximal to distal, up to 2.0 *cun*. {Six Bowels and Lung Auxiliary Reaction Areas} (Jian¹ Zhong¹ is located 2.5 *cun* distal to LI-15^{WHO} on the Hand Yang² Ming² Large Intestine Channel.)
- **Wu³ Ling³:** These points are found on five rostral-caudal lines located on the back and they should be ‘let’ rather than needled with filiform needles. The first row of 10 points is located on the Du Mai (Governing Vessel) in the interspinous spaces between: T₂/T₃, T₃/T₄, T₄/T₅, T₅/T₆, T₆/T₇, T₇/T₈, T₈/T₉, T₉/T₁₀, T₁₀/T₁₁, and T₁₁/T₁₂. The second row of 8 points (per row) is located 3.0 *cun* lateral on either side of the Du Mai (Governing Vessel), commences from the level of the T₂/T₃, and includes points at the levels of T₃/T₄, T₄/T₅, T₅/T₆, T₆/T₇, T₇/T₈, T₈/T₉, T₉/T₁₀. The third row of 7 points (per row) is located 6.0 *cun* lateral on either side of the Du Mai (Governing Vessel), commences from the level of the T₂/T₃, and includes points at the levels of T₃/T₄, T₄/T₅, T₅/T₆, T₆/T₇, T₇/T₈, and T₈/T₉. (In this case, use those points on the first and second rows.)
- **San¹ Jiang¹:** This group of points is located on the dorsal surface of the lower back as follows: The six midsagittal points are located along the vertebral column, in the interspinous spaces between L₁/L₂, L₂/L₃, L₃/L₄, L₄/L₅, L₅/S₁, and S₁/S₂; the lateral points are located 3.0 *cun* lateral to the midsagittal plane on either side of the aforementioned interspinous levels; ‘Let’ with a prismatic needle. {Results will be improved if one can ‘let’ the ‘Malign Blood’, then needle Shen⁴ Guan¹ – i.e., 2.5 *cun* distal to Sp-9^{WHO}}

Pictures are on page 261; scroll down on this pdf

Paresthesiae: Pedal (LKC-2)



‘Let’ Wu³ Ling³ NO PNEUMOTHORAX



‘Let’ San¹ Jiang¹



‘Let’ Affected Foot